

Laser cooling all the way down to molecular condensate

J. Dziarmaga¹ and M. Lewenstein²

¹ *Institute of Physics and Centre for Complex Systems, Jagiellonian University, ul. Reymonta 4, 30-059 Kraków, Poland*

² *Institut für Theoretische Physik, Universität Hannover, D-30167 Hannover, Germany*

(January 25, 2005)

Numerical simulations show that laser cooling of fermions on the repulsive side of the Feshbach resonance can sympathetically cool molecules well below their condensation temperature.

PACS: 03.75.Ss, 39.25.+k, 03.75-b

Fermi superfluids are in the center of interest of recent studies in physics of ultracold quantum gases. They provide perhaps the most promising model system to study the superfluidity, and the Bardeen-Cooper-Schrieffer (BCS) theory in the limit of strong interactions, when the size of Cooper pairs becomes comparable or even smaller than average interparticle distance, and the mean field description breaks down [1]. In cold atom systems the use of Feshbach resonances permits to tune the s -wave scattering length between the atoms with opposite “spin” [2], assuring in this way a high flexibility of the systems. This technique allows in particular to drive the system through the BCS-BEC crossover when the scattering length diverges and changes sign from negative (attractive) to positive (repulsive). During this passage, the state of the system continuously evolves from the BCS superfluid of weakly correlated Cooper pairs, through the pseudogap regime, and then the unitarity limit, when the scattering length diverges, to the Bose-Einstein condensate (BEC) of diatomic molecules. By now several successful experiments with molecular condensates were performed [3] which opened a way to first successful experiments with Fermi superfluids [4].

Earlier theoretical studies demonstrated that laser cooling allows to reach the degeneracy in Fermi systems [5], and to go all the way down to the Fermi superfluid [6] on the attractive side of the Feshbach resonance. The purpose of this Communication is to show that laser cooling offers also an alternative technique to reach molecular condensate on the repulsive side of the resonance. Moreover, it allows to post-cool the condensate prepared via evaporative cooling down to even lower temperatures. We have developed here a fully self-consistent quantum kinetic theory of molecular condensate creation via laser cooling¹. The results are very promising: despite the fact that condensate formation shifts the energies of quasi-particle excitations with respect to the equally spaced harmonic oscillator levels, and at the same time reduces

fermionic population exposed to laser cooling, the cooling remains efficient well below the condensation temperature of $0.18T_F$ down to at least $0.06T_F$. Optimization of the cooling protocol could probably significantly reduce this temperature.

We consider system containing two species of fermions with “spin up and down”, molecules which are in their bound states, and fermions in an excited state. We work on the repulsive side of the Feshbach resonance, where interactions strengths between spin up/down fermions and molecules, as well as the binding energy of the molecules are simple universal functions of the effective s -wave scattering length a between spin up and down fermions [8]. Close to the resonance the binding energy of molecules $\nu = -\frac{\hbar^2}{ma^2}$ can be made less than the condensation temperature $k_B T_C$ for the molecules. As a result at all temperatures greater than T_C populations of molecules and fermions are comparable. Below T_C , a fraction of molecules condenses leaving room in molecular phase space for more fermions to bind into molecules. Fermionic population is then shrinking, but it remains still comparable to the shrinking population of non-condensed molecules all the way down to the dissociation temperature, $T_D = \nu/k_B$. Below T_D , fermionic population is suppressed, but at the same time almost all molecules are already condensed. Thus, close to the Feshbach resonance, when $T_D < T_C$, there are favorable conditions for efficient sympathetic cooling of molecules by collisions with spin up/down fermions at all temperatures of interest. The fermions in turn are subject to laser cooling, and provide the heat reservoir, which is comparable to the cooled system (non-condensed molecules). Populations of the spin up and down fermions are the same. Mutual s -wave interactions between these two populations and the population of molecules lead to thermalization on a time scale, which is faster than the rate of laser cooling. At the same time three-body collisions keep the system close to chemical equilibrium between molecules and fermions. The system remains thus in a state of quasi-equilibrium with slowly decreasing temperature. This thermal state is described by the generalized BCS theory or boson-fermion model [9] which includes BEC of molecules described by the Gross-Pitaevskii equation. The model takes into account coherent tunneling between molecules and pairs of spin up and down fermions. Thermal excitations of the molecular condensate are described by the bosonic Bogoliubov theory. Structure of the ground state of fermions and its fermionic excitations

¹similar to self-consistent theories of the BEC growth [7]

follows from the set of Bogoliubov-de Gennes equations together with self-consistent definitions of the gap function and the mean field potential. On the repulsive side of the Feshbach resonance the leading contribution to the gap function comes from the molecular BEC through the coherent coupling between molecules and pairs of fermions. In particular the gap function experienced by fermions is proportional to the condensate wave function. Laser cooling of fermions was described in detail in Ref. [5]. Coherent laser excites atoms from, say, the spin down ground state to the excited state, and spontaneous emission brings them back to the spin down ground state. Frequencies of the laser assure that the generalized Raman cooling takes place [10]. In this Communication we take into account interactions between fermions and describe the cooling process in terms of instantaneous fermionic and bosonic (molecular) Bogoliubov quasiparticles, whose eigenfunctions and eigenenergies are self-consistently updated during the evolution. We work in the “festina lente” limit to avoid reabsorption effects [11], and also employ spherical symmetry and ergodic approximations [6,7].

The boson-fermion model [9] is defined by the Hamiltonian

$$\hat{H} = \int d^3r \left[\sum_{a=+, -, e, m} \hat{\psi}_a^\dagger \mathcal{H}_a \hat{\psi}_a + (\nu - \mu) \hat{\psi}_m^\dagger \hat{\psi}_m + \lambda \hat{\psi}_m^\dagger \hat{\psi}_+ \hat{\psi}_- + \text{h.c.} + g \hat{\psi}_+^\dagger \hat{\psi}_- \hat{\psi}_- \hat{\psi}_+ + 0.3 g \hat{\psi}_m^\dagger \hat{\psi}_m^\dagger \hat{\psi}_m \hat{\psi}_m + 1.2 g \hat{\psi}_m^\dagger \hat{\psi}_m \left(\hat{\psi}_+^\dagger \hat{\psi}_+ + \hat{\psi}_-^\dagger \hat{\psi}_- \right) \right], \quad (1)$$

where the fields $\hat{\psi}_+$ and $\hat{\psi}_-$ describe fermions with spin up and down, respectively, $\hat{\psi}_e$ corresponds to fermions in the excited state, and $\hat{\psi}_m$ is the bosonic molecular field. Here $\mathcal{H}_a = -\frac{\hbar^2}{2m_a} \nabla^2 + V_a(\vec{r}) - \mu$ is a single particle Hamiltonian, m_a is atomic mass m or molecular mass $2m$, $V_a(\vec{r}) = \frac{1}{2} m_a \Omega_a^2 r^2$ is the harmonic trap potential, μ is chemical potential, $g = \frac{4\pi\hbar^2 a}{m}$ is fermion-fermion interaction strength with a large positive s -wave scattering length a , λ is a coupling between pairs of atoms and molecules, and $\nu = -\frac{\hbar^2}{ma^2}$ is the molecular binding energy. Close to the Feshbach resonance, when g is large, we can neglect interactions with excited atoms.

In the mean field approximation, which closely follows the BCS theory, the quartic fermion-fermion interaction term in the Hamiltonian (1) is made quadratic by replacing products of operators by their averages in all possible ways. The averages are the mean field potential $W(\vec{r}) = 1.2 g \langle \hat{\psi}_m^\dagger(\vec{r}) \hat{\psi}_m(\vec{r}) \rangle + g \langle \hat{\psi}_\pm^\dagger(\vec{r}) \hat{\psi}_\pm(\vec{r}) \rangle$, and the anomalous pairing potential $P(\vec{r}) = \langle \hat{\psi}_+(\vec{r}) \hat{\psi}_-(\vec{r}) \rangle$, which is mixing fermions with spin up and down. In the mean-field approximation the molecular field $\hat{\psi}_m$ is

replaced by a c -number condensate amplitude ϕ that fulfills the stationary Gross-Pitaevskii equation

$$(2\mu - \nu)\phi = \mathcal{H}_m \phi + 0.6 g |\phi|^2 \phi + \lambda P. \quad (2)$$

Here ϕ is normalized to the number of condensed molecules. The mean-field equations for fermions are

$$i\hbar \frac{d}{dt} \hat{\psi}_\pm = \mathcal{H}_\pm \hat{\psi}_\pm + W(\vec{r}) \hat{\psi}_\pm \mp \Delta(\vec{r}) \hat{\psi}_\mp^\dagger, \quad (3)$$

with an effective gap function $\Delta(\vec{r}) = gP(\vec{r}) + \lambda\phi(\vec{r})$. These equations are mixing spin up and down components but they can be “diagonalized” by the Bogoliubov transformation $\hat{\psi}_\pm(\vec{r}) = \sum_m \hat{b}_{m,\pm} u_m(\vec{r}) \mp \hat{b}_{m,\mp}^\dagger v_m^*(\vec{r})$, with fermionic quasiparticle annihilation operators $\hat{b}_{m,\pm}$. The quasiparticle modes (u_m, v_m) fulfill the Bogoliubov-de Gennes equations with positive energies ω_m ,

$$\begin{aligned} \omega_m u_m &= + \mathcal{H}_\pm u_m + W u_m - \Delta v_m, \\ \omega_m v_m &= - \mathcal{H}_\pm v_m - W v_m - \Delta^* u_m. \end{aligned} \quad (4)$$

In a thermal state with inverse temperature β the average occupation numbers of quasiparticle states are $N_m = \langle \hat{b}_{m,\pm}^\dagger \hat{b}_{m,\pm} \rangle = [\exp(\beta\omega_m) + 1]^{-1}$. Equations (4) are solved together with the self-consistency conditions

$$\Delta(\vec{r}) = \lambda\phi(\vec{r}) + g \sum_m [1 - 2N_m] u_m(\vec{r}) v_m^*(\vec{r}), \quad (5)$$

$$\begin{aligned} W(\vec{r}) &= 1.2 g \langle \hat{\psi}_m^\dagger(\vec{r}) \hat{\psi}_m(\vec{r}) \rangle + \\ &g \sum_m (1 - N_m) |v_m(\vec{r})|^2 + N_m |u_m(\vec{r})|^2, \end{aligned} \quad (6)$$

by successive iterations with the chemical potential μ adjusted to keep the total number of atoms constant. The average molecular density $\langle \hat{\psi}_m^\dagger(\vec{r}) \hat{\psi}_m(\vec{r}) \rangle$ is a sum of the condensate density $|\phi(\vec{r})|^2$ plus density of non-condensed molecules. After every iteration the Gross-Pitaevskii equation (2) is solved by relaxation. The total number of atoms includes atoms bound into non-condensed molecules depleted from the condensate by thermal and quantum fluctuations. We estimate the number of non-condensed molecules using the approximate bosonic Bogoliubov modes in the Thomas-Fermi limit [12]. The ultraviolet divergence in Eq.(5) is regularized by the quickly convergent method of Ref. [13].

Excitation of atoms from the spin down state to the excited state is described by the Hamiltonian

$$\hat{H}_{\text{las}} = \int d^3r \left[\frac{1}{2} \Omega e^{i\vec{k}_L \cdot \vec{r}} \hat{\psi}_e^\dagger(\vec{r}) \hat{\psi}_-(\vec{r}) + \text{h.c.} - \delta \hat{\psi}_e^\dagger \hat{\psi}_e \right], \quad (7)$$

driving coherent oscillations with Rabi frequency Ω and laser detuning δ . The excitation is accompanied by the spontaneous emission described by a superoperator

$$\mathcal{L} \hat{\rho} = \gamma \sum_{mk} U_{mk} \mathcal{D}[\hat{b}_{m,-}^\dagger \hat{e}_k] \rho + V_{mk} \mathcal{D}[\hat{b}_{m,+} \hat{e}_k] \rho \quad (8)$$

with a spontaneous emission rate γ . Here, the Lindblad superoperator is $\mathcal{D}[\hat{A}] \rho = \hat{A} \rho \hat{A}^\dagger - \frac{1}{2} \hat{A}^\dagger \hat{A} \rho - \frac{1}{2} \hat{A} \hat{A}^\dagger \rho$, and the matrix elements are e.g. $U_{mk} = \int d\Omega_k \mathcal{W}(\Omega_k) |u_{mk}(\vec{k})|^2$ with the spontaneous emission pattern $\mathcal{W}(\Omega_k)$ and the generalized Frank-Condon factors $u_{mk}(\vec{k}) = \int d^3r e^{i\vec{k}\vec{r}} w_k^*(\vec{r}) u_m(\vec{r})$. Here, $w_k(\vec{r})$ is the k -th eigenstate of the harmonic oscillator and $\hat{e}_k$ is an annihilation operator of an excited atom in this state.

Adiabatic elimination of the excited state (cf. [14]) leads to the kinetic equations for the occupation numbers taking into account the effects of Fermi statistics

$$\begin{aligned} \frac{dN_{m,-}}{dt} &= \sum_n \Gamma_{m \leftarrow n}^{(-)} (1 - N_{m,-}) N_{n,-} - (n \leftrightarrow m) + \\ &\quad C_{mn} (1 - N_{m,-}) (1 - N_{n,+}) - A_{nm} N_{m,-} N_{n,+}, \\ \frac{dN_{m,+}}{dt} &= \sum_n \Gamma_{m \leftarrow n}^{(+)} (1 - N_{m,+}) N_{n,+} - (n \leftrightarrow m) + \\ &\quad C_{nm} (1 - N_{m,+}) (1 - N_{n,-}) - A_{mn} N_{m,+} N_{n,-}. \end{aligned}$$

They describe relaxation of $\pm$ quasiparticles ($\Gamma_{m \leftarrow n}^{(\pm)}$), and creation/annihilation of pairs of $+$ and $-$ quasiparticles (C_{nm} and A_{nm}). The transition rates are e.g.

$$\Gamma_{m \leftarrow n}^{(-)} = \frac{\Omega^2}{2\gamma} \sum_k \frac{\gamma^2 U_{mk} |u_{nk}(\vec{k}_L)|^2}{(\delta - \omega_k^e + \omega_n)^2 + \gamma_k^2}. \quad (9)$$

Here, ω_k^e is the energy of the k -th harmonic oscillator state and γ_k is approximate spontaneous decay rate of an excited atom in this state, $\gamma_k = \gamma \sum_m U_{mk} (1 - N_{m,-}) + V_{mk} N_{m,+}$, see Appendix A.

Laser cooling drives average occupation numbers $N_{m,\pm}$ of fermionic quasiparticles out of thermal equilibrium. At the same time interactions between fermionic and bosonic quasiparticles drive the system towards thermal equilibrium. Numerical simulations in Ref. [5] show that thermal relaxation remains efficient all the way down to the temperature of $0.03 T_F$. This justifies our assumption of fast equilibration to a quasi-equilibrium state. After a short period of laser cooling dt the occupation numbers $N_{m,\pm}$ go out of equilibrium where $N_{m,\pm} = N_m = [\exp(\beta\omega_m) + 1]^{-1}$. The initial total energy $E(T) = \sum_{m,\pm} \omega_m N_{m,\pm} + \sum_m \omega_m^B N_m^B$, including bosonic Bogoliubov quasiparticles with frequencies ω_m^B , and average occupation numbers $N_m^B = [\exp(\beta\omega_m^B) - 1]^{-1}$, changes by $dE = \sum_m \omega_m (N_{m,+} + N_{m,-} - 2N_m)$. The relaxation brings the occupation numbers $N_{m,\pm}$ to a new state of equilibrium at a temperature $T + dT$, but does not change the total energy. The energy of the system at the new temperature $E(T + dT)$ differs from the initial $E(T)$ by $dE_T \approx 2 \frac{dT}{T^2} \sum_m \omega_m^2 N_m (1 - N_m) + \frac{dT}{T^2} \sum_m (\omega_m^B)^2 N_m^B (1 + N_m^B)$. Conservation of energy in thermal relaxation means that $dE = dE_T$. In our

simulations we use this equality to find the new lower temperature $T + dT$, and then solve the Bogoliubov-de Gennes equations and the Gross-Pitaevskii equation self-consistently to adjust the Bogoliubov modes, the condensate wave function and the chemical potential to the new lower temperature. With the new Bogoliubov modes we calculate new transition rates, etc.

Calculation of the transition rates is the most time consuming part of the numerical simulation. This is why we were forced to assume spherical symmetry. For spherically symmetric $\Delta(r)$, $W(r)$ and $\phi(r)$ the Bogoliubov modes are $u_{ln}(r) Y_{lm}(\theta, \varphi)$ and their energies ω_{ln} do not depend on m . The bosonic quasiparticle energies in the Thomas-Fermi limit [12] become $\omega_{ln}^B = \Omega_m (2n^2 + 2nl + 3n + l)^{1/2}$. To be consistent with the spherical symmetry we also made ergodic approximation that quasiparticle occupation numbers $N_{ln,\pm}$ do not depend on m . In other words, we assume fast equilibration within each quasiparticle energy shell which keeps the system in a spherically symmetric state. More on spherical symmetry in Appendix B.

In our simulations we use 81 harmonic oscillator levels, and assume that there are $N = 10660$ atoms with spin up and down which in the non-interacting case at $T = 0$ is equivalent to 39 filled energy levels. The frequency of the isotropic trap is $\Omega_a = 2\pi \cdot 2400 \text{ Hz}$, the same for all fermions and molecules. The scattering length for the interaction between the two species is $a = 1200 \text{ \AA}$, a realistic value for interactions between two spin states $|F = 9/2, m_F = 9/2\rangle$ and $|F = 9/2, m_F = 7/2\rangle$ of ^{40}K near the Feshbach resonance. In the natural harmonic oscillator units this scattering length gives the interaction strength $g = 4.7$, and the molecular binding energy $\nu = -7.3$. The wavelength of the cooling laser is $\lambda = 720 \text{ nm}$, and the laser detuning is $\delta = -12\hbar\Omega_a$. In all simulations the frequency of Rabi oscillations is much less than the spontaneous emission rate, $\Omega = 0.1\gamma$, so that the average occupation of excited state remains small. In our simulations we neglect $W(r)$. The main effect of the mean field potential is an overall shift of energy levels which can be compensated by laser detuning δ . The coherent tunneling rate in trap units is $\lambda = 1$.

In Fig. 1 we show temperature as a function of time for laser cooling with two values of γ . Similarly as on the BCS side of the Feshbach resonance the nonzero gap function $\Delta(r)$ induced by the molecular BEC does not affect much the efficiency of laser cooling. Moreover, the cooling remains efficient even below the dissociation temperature of $0.18 T_F$. This is not quite surprising because even at such low temperature there is a non-zero fraction of fermions induced by coherent tunneling from the molecular BEC, which can be cooled by the laser. The initial cooling is faster for $\gamma = 10$ but the final temperature of $0.06 T_F$ is lower for $\gamma = 1.25$. The best strategy to reach low temperatures in reasonable time is to do the cooling in a few stages starting with large γ and finishing

with small γ , as shown in the inset in Fig.1. In Fig. 2 we show the growth of the molecular condensate in the same simulations as in Fig.1.

To summarize, our numerical simulations show that for reasonable experimental parameters consistent with the requirements of the Festina lente regime, it is possible to laser cool fermions on the positive side of the Feshbach resonance all the way down to the molecular BEC. Despite the pairing effects, which produce a gap in the quasiparticle energy spectrum and the heat capacity of non-condensed molecules, it is possible to reach at least $0.06 T_F$ in a time of a few seconds.

We would like to thank M. Baranov, L. Santos, and Z. Idziaszek for discussions. This work was supported in part by ESF QUEDDIS program, DFG (SFB 407, SPP1116, 432 POL), and the KBN grant PBZ-MIN-008/P03/2003.

-
- [1] A.J. Leggett, in *Modern Trends in the Theory of Condensed Matter*, A. Pekalski and R. Przystawa (Eds.), (Springer, Berlin, 1980); J. Phys. (Paris) Colloq. **41**, c7-19 (1980); P. Nozières and S.Schmitt-Rink, J. Low Temp. Phys. **59**, 195 (1985); M. Randeria, in *Bose-Einstein Condensation*, A. Griffin, D. Snoke and S. Stringari (Eds.), (Cambridge University Press, Cambridge 1995).
 - [2] E. Timmermans, P. Tommasini, M. Hussein, and A. Kerman, Phys. Rep. **315**, 199 (1999).
 - [3] S. Jochim *et al.*, Denschlag, and R. Grimm, Science **302**, 2101 (2003); C.A. Regal, M. Greiner, and D.S. Jin, Phys. Rev. Lett. **92**, 040403 (2004); M.W. Zwierlein *et al.*, Phys. Rev. Lett. **92**, 120403 (2004); J. Kinast *et al.*, Phys. Rev. Lett. **92**, 150402 (2004); T. Bourdel *et al.*, L. Tarruell, S.J.J.M.F. Kokkelmans, Phys. Rev. Lett.**93**, 050401 (2004).
 - [4] C. Chin *et al.*, and R. Grimm, Science **305**, 1128 (2004); for theory see J. Kinnunen, M. Rodriguez, and P. Törmä, Phys. Rev. Lett. **92**, 120403 (2004); Science **305**, 1131 (2004); M. Greiner, C.A. Regal, and D. Jin, cond-mat/0407381; M.W. Zwierlein *et al.* cond-mat/0412675; J. Zheng *et al.* cond-mat/0410167; M. Bartenstein *et al.*, cond-mat/0412712.
 - [5] Z. Idziaszek, L. Santos, M. Baranov, and M. Lewenstein, Phys. Rev. A**67**, 041403 (2003); Z. Idziaszek, L. Santos, M. Baranov, and M. Lewenstein, J. Opt. B**5**, 190 (2003).
 - [6] J. Dziarmaga and M. Lewenstein, cond-mat/0410704, Phys.Rev.Lett. **94**, XXXXX (2005).
 - [7] D. Jaksch, C.W. Gardiner, and P. Zoller, Phys. Rev. A**56**, 575 (1997); D. Jaksch, C.W. Gardiner K.M. Gheri, and P. Zoller, *ibid.* A**58**, 1450 (1998); C.W. Gardiner and P. Zoller, *ibid.*A**61**, 033601 (2000).
 - [8] D.S. Petrov, C. Salomon and G.V. Shlyapnikov, Phys. Rev. Lett. **93**, 090404 (2004); Phys. Rev. A **71**, 012708 (2005).
 - [9] H.T.C. Stoof, M. Houbiers, C.A. Sackett, and

- R.G. Hulet, Phys. Rev. Lett.**76**, 10 (1996); E. Timmermans, K. Furuya, P.W. Milonni, and A.K. Kerman, Phys. Lett. A **285**, 228 (2001); M. Holland, S.J.J.M.F. Kokkelmans, M.L. Chiofalo, and R. Walser, Phys. Rev. Lett. **87**, 120406 (2001); S.J.J.M.F. Kokkelmans, G.V. Shlyapnikov, and C. Salomon, cond-mat/0308384.
- [10] L. Santos, Z. Idziaszek, J.I. Cirac, and M. Lewenstein, J. Phys. B**33**, 4131 (2000).
- [11] Y. Castin, J.I. Cirac, and M. Lewenstein, Phys. Rev. Lett. **80**, 5305 (1998).
- [12] S. Stringari, Phys.Rev.Lett.**77**, 2360 (1996).
- [13] M. Grasso and M. Urban, Phys. Rev. A**68**, 033610 (2003).
- [14] L. Santos and M. Lewenstein, Phys. Rev. A**60**, 3851 (1999); Eur. Phys. J. D**7**, 379 (1999).
- [15] For more details see J. Dziarmaga and M. Lewenstein,




FIG. 1. Temperature T/T_F as a function of time for $\gamma = 10$ (solid line) and $\gamma = 1.25$ (dashed line). Here $T_F = 40.5\hbar\Omega_a$, and the observed $T_C \approx 0.6T_F$. The cooling is faster for $\gamma = 10$, but the lower final temperature of $0.06T_F$ is achieved for $\gamma = 1.25$. The inset shows T/T_F in a two stage cooling




FIG. 2. The number of condensed molecules in the same simulations as in Fig. 1.

APPENDIX A: TRANSITION RATES

Laser cooling is realized by the Hamiltonian

$$\hat{H}_{las} = \frac{1}{2}\Omega \int d^3r e^{i\vec{k}_L \vec{r}} \hat{\psi}_e^\dagger(\vec{r}) \hat{\psi}_-(\vec{r}) + \text{h.c.} , \quad (10)$$

driving coherent oscillations with Rabi frequency Ω between the excited state e and the state $-$, together with the spontaneous emission superoperator

$$\begin{aligned} \mathcal{L} \hat{\rho} = & \gamma \int d\varphi d\cos\theta \mathcal{W}(\theta, \varphi) \int d^3r_1 d^3r_2 e^{i\vec{k}(\vec{r}_1 - \vec{r}_2)} \\ & \times \left[2\hat{\psi}_-^\dagger(\vec{r}_2) \hat{\psi}_e(\vec{r}_2) \hat{\rho} \hat{\psi}_e^\dagger(\vec{r}_1) \hat{\psi}_-(\vec{r}_1) - \hat{\psi}_e^\dagger(\vec{r}_1) \hat{\psi}_-(\vec{r}_1) \hat{\psi}_-^\dagger(\vec{r}_2) \hat{\psi}_e(\vec{r}_2) \hat{\rho} - \hat{\rho} \hat{\psi}_e^\dagger(\vec{r}_1) \hat{\psi}_-(\vec{r}_1) \hat{\psi}_-^\dagger(\vec{r}_2) \hat{\psi}_e(\vec{r}_2) \right] , \end{aligned} \quad (11)$$

with an effective spontaneous emission rate of 2γ and the fluorescence dipole pattern $\mathcal{W}(\theta, \varphi)$. Expansion in the eigenmodes $w_l(\vec{r})$ of the harmonic trap

$$\hat{\psi}_e(\vec{r}) = \sum_l \hat{e}_l w_l(\vec{r}) , \quad (12)$$

and the Bogoliubov transformation (see main text) give

$$\hat{H}_{las} = \frac{1}{2}\Omega \sum_{ml} \left[\hat{b}_{m,-} u_{ml}(\vec{k}_L) + \hat{b}_{m,+}^* v_{ml}^*(\vec{k}_L) \right] \hat{e}_l^\dagger + \text{h.c.} . \quad (13)$$

Here the (generalized) Frank-Condon factors are

$$\begin{aligned} u_{ml}(\vec{k}) &= \int d^3r e^{i\vec{k}\vec{r}} w_l^*(\vec{r}) u_m(\vec{r}) , \\ v_{ml}^*(\vec{k}) &= \int d^3r e^{i\vec{k}\vec{r}} w_l^*(\vec{r}) v_m^*(\vec{r}) . \end{aligned} \quad (14)$$

In a similar way, and after rotating wave approximation for e -atoms and the Bogoliubov quasiparticles, the spontaneous emission term becomes

$$\begin{aligned} \mathcal{L} \hat{\rho} = & \gamma \sum_{ml} U_{ml} \left(2\hat{b}_{m,-}^\dagger \hat{e}_l \hat{\rho} \hat{e}_l^\dagger \hat{b}_{m,-} - \hat{e}_l^\dagger \hat{e}_l \hat{b}_{m,-} \hat{b}_{m,-}^\dagger \hat{\rho} - \hat{\rho} \hat{e}_l^\dagger \hat{e}_l \hat{b}_{m,-} \hat{b}_{m,-}^\dagger \right) + \\ & \gamma \sum_{ml} V_{ml} \left(2\hat{b}_{m,+}^\dagger \hat{e}_l \hat{\rho} \hat{e}_l^\dagger \hat{b}_{m,+} - \hat{e}_l^\dagger \hat{e}_l \hat{b}_{m,+} \hat{b}_{m,+}^\dagger \hat{\rho} - \hat{\rho} \hat{e}_l^\dagger \hat{e}_l \hat{b}_{m,+} \hat{b}_{m,+}^\dagger \right) . \end{aligned} \quad (15)$$

Here the spontaneous emission rates are

$$\begin{aligned} U_{ml} &= \int d\varphi d\cos\theta \mathcal{W}(\theta, \varphi) |u_{ml}(\vec{k})|^2 , \\ V_{ml} &= \int d\varphi d\cos\theta \mathcal{W}(\theta, \varphi) |v_{ml}(\vec{k})|^2 . \end{aligned} \quad (16)$$

Adiabatic elimination of the excited state e , similar as in Ref. [14], results in the following evolution equations for the occupation numbers

$$\frac{dN_{m,-}}{dt} = \sum_n \Gamma_{m \leftarrow n}^{(-)} (1 - N_{m,-}) N_{n,-} - \Gamma_{n \leftarrow m}^{(-)} (1 - N_{n,-}) N_{m,-} + C_{mn} (1 - N_{m,-}) (1 - N_{n,+}) - A_{nm} N_{m,-} N_{n,+} , \quad (17)$$

$$\frac{dN_{m,+}}{dt} = \sum_n \Gamma_{m \leftarrow n}^{(+)} (1 - N_{m,+}) N_{n,+} - \Gamma_{n \leftarrow m}^{(+)} (1 - N_{n,+}) N_{m,+} + C_{nm} (1 - N_{m,+}) (1 - N_{n,-}) - A_{mn} N_{m,+} N_{n,-} . \quad (18)$$

There are contributions from four different processes:

- relaxation of $-$ quasiparticles

$$\Gamma_{m \leftarrow n}^{(-)} = \frac{\Omega^2}{2\gamma} \sum_l \frac{\gamma^2 U_{ml} |u_{nl}(\vec{k}_L)|^2}{|\delta - \omega_l^e + \omega_n + i\gamma \sum_s U_{sl}(1 - N_{s,-} + \delta_{s,n}) + V_{sl} N_{s,+}|^2}, \quad (19)$$

- relaxation of $+$ quasiparticles

$$\Gamma_{m \leftarrow n}^{(+)} = \frac{\Omega^2}{2\gamma} \sum_l \frac{\gamma^2 V_{nl} |v_{ml}(\vec{k}_L)|^2}{|\delta - \omega_l^e - \omega_m + i\gamma \sum_s U_{sl}(1 - N_{s,-}) + V_{sl}(N_{s,+} + \delta_{m,s})|^2}, \quad (20)$$

- creation of pairs of $+$ and $-$ quasiparticles

$$C_{mn} = \frac{\Omega^2}{2\gamma} \sum_l \frac{\gamma^2 U_{ml} |v_{nl}(\vec{k}_L)|^2}{|\delta - \omega_l^e - \omega_n + i\gamma \sum_s U_{sl}(1 - N_{s,-}) + V_{sl}(N_{s,+} + \delta_{n,s})|^2}, \quad (21)$$

- annihilation of pairs of $+$ and $-$ quasiparticles

$$A_{mn} = \frac{\Omega^2}{2\gamma} \sum_l \frac{\gamma^2 V_{ml} |u_{nl}(\vec{k}_L)|^2}{|\delta - \omega_l^e + \omega_n + i\gamma \sum_s U_{sl}(1 - N_{s,-} + \delta_{n,s}) + V_{sl} N_{s,+}|^2}. \quad (22)$$

APPENDIX B: SPHERICAL SYMMETRY

Simulation of laser cooling with a non-vanishing pairing function $P(\vec{r})$, and a Hartree potential $g_0 \rho(\vec{r})$ is a numerically hard problem. The main difficulty is that after every short period of laser cooling it is necessary to reiterate Bogoliubov-de Gennes equations in order to obtain a new self-consistent pairing function, Hartree potential, and the Bogoliubov modes (ω_m, u_m, v_m) . With the new Bogoliubov modes, the transition rates $A_{mn}, C_{mn}, \Gamma_{m \leftarrow n}^{\pm}$ have to be calculated again, and this is a very time-consuming operation. This numerical effort is reduced a lot with assumption of spherical symmetry. With spherically symmetric $\Delta(r)$ and $\rho(r)$ the Bogoliubov modes can be decomposed into spherical and radial parts

$$u_{ln}(r) Y_{lm}(\theta, \varphi), \quad (23)$$

$$v_{ln}(r) Y_{lm}(\theta, \varphi). \quad (24)$$

In a similar way harmonic oscillator modes become $W_{ln}(r) Y_{lm}(\theta, \varphi)$. Quasiparticle energies ω_{ln} do not depend on m . We assume fast equilibration within each degenerate energy shell nl so that all occupation numbers in a given shell are the same $N_{ln,\pm}$. This assumption greatly simplifies the rate equations

$$\begin{aligned} \frac{d}{dt} N_{l_1 n_1, -} = & \sum_{l_2 n_2} \gamma_{l_1 n_1 \leftarrow l_2 n_2}^{(-)} (1 - N_{l_1 n_1, -}) N_{l_2 n_2, -} - \gamma_{l_2 n_2 \leftarrow l_1 n_1}^{(-)} (1 - N_{l_2 n_2, -}) N_{l_1 n_1, -} + \\ & c_{l_1 n_1 l_2 n_2} (1 - N_{l_1 n_1, -}) (1 - N_{l_2 n_2, +}) - a_{l_2 n_2 l_1 n_1} N_{l_1 n_1, -} N_{l_2 n_2, +}, \end{aligned} \quad (25)$$

$$\begin{aligned} \frac{d}{dt} N_{l_1 n_1, +} = & \sum_{l_2 n_2} \gamma_{l_1 n_1 \leftarrow l_2 n_2}^{(+)} (1 - N_{l_1 n_1, +}) N_{l_2 n_2, +} - \gamma_{l_2 n_2 \leftarrow l_1 n_1}^{(+)} (1 - N_{l_2 n_2, +}) N_{l_1 n_1, +} + \\ & c_{l_2 n_2 l_1 n_1} (1 - N_{l_1 n_1, +}) (1 - N_{l_2 n_2, -}) - a_{l_1 n_1 l_2 n_2} N_{l_1 n_1, +} N_{l_2 n_2, -}. \end{aligned} \quad (26)$$

Here $\gamma^{(\pm)}, a, c$ are averaged transition rates, for example

$$\gamma_{l_1 n_1 \leftarrow l_2 n_2}^{(-)} = \frac{1}{2l_1 + 1} \sum_{m_1 m_2} \Gamma_{l_1 n_1 m_1 \leftarrow l_2 n_2 m_2}^{(-)}. \quad (27)$$

This equation combined together with the definition of $\Gamma^{(-)}$ gives

$$\gamma_{l_1 n_1 \leftarrow l_2 n_2}^{(-)} = \frac{\gamma \Omega^2}{2(2l_1 + 1)} \sum_{m_1 m_2} \sum_{l_e n_e m_e} \frac{U_{l_1 n_1 m_1 l_e n_e m_e} |u_{l_2 n_2 m_2 l_e n_e m_e}(\vec{k}_L)|^2}{[\delta - \omega(l_e + 2n_e - \mu) + \omega_{l_2 n_2}]^2 + \gamma^2 R_{l_e n_e}^2}. \quad (28)$$

Here the frequency of the intermediate excited state $\omega(l_e + 2n_e - \mu)$ is measured with respect to the chemical potential μ because the quasiparticle energies $\omega_{l_2 n_2}$ are also defined with respect to μ . For a laser beam with a $\vec{k}_L$ along the $\hat{z}$ -axis the indices m_e and m_2 must be the same. Furthermore, it follows from the properties of Clebsch-Gordan coefficients that the sum $\sum_{m_1} U_{l_1 n_1 m_1 l_e n_e m_e}$ does not depend on m_e so that we can set e.g. $m_e = 0$ and

$$\begin{aligned} \gamma_{l_1 n_1 \leftarrow l_2 n_2}^{(-)} &= \frac{\gamma \Omega^2}{2(2l_1 + 1)} \sum_{l_e n_e} \frac{(\sum_{m_1} U_{l_1 n_1 m_1 l_e n_e 0}) \left(\sum_{m_2} |u_{l_2 n_2 m_2 l_e n_e m_2}(\vec{k}_L)|^2 \right)}{[\delta - \omega(l_e + 2n_e - \mu) + \omega_n]^2 + \gamma^2 R_{l_e n_e}^2} \equiv \\ &= \frac{\gamma \Omega^2}{2(2l_1 + 1)} \sum_{l_e n_e} \frac{SU_{l_1 n_1 l_e n_e} \quad su_{l_2 n_2 l_e n_e}}{[\delta - \omega(l_e + 2n_e - \mu) + \omega_n]^2 + \gamma^2 R_{l_e n_e}^2}. \end{aligned} \quad (29)$$

The last form shows that the transition rate $\gamma^{(-)}$ can be constructed out of matrices SU , su and a vector R . These elements can be expressed through even more elementary building blocks

$$\begin{aligned} SU_{l_1 n_1 l_e n_e} &= \sum_l (jWu)_{l_e n_e l_1 n_1}^2 S_{l_e l_1}, \\ su_{l_2 n_2 l_e n_e} &= \sum_m \left| \sum_l s_{l_e l_2 m} (jWu)_{l_e n_e l_2 n_2} \right|^2. \end{aligned}$$

The more elementary building blocks are

$$\begin{aligned} (jWu)_{l_e n_e l_1 n_1} &= \int r^2 dr j_l(kr) W_{l_e n_e}(r) u_{l_1 n_1}(r), \\ S_{l_e l_1} &= (2l + 1) \frac{2l_e + 1}{2l_1 + 1} \sum_m \langle l, l_e, m, 0 | l_1, m \rangle^2 \langle l, l_e, 0, 0 | l_1, 0 \rangle^2, \\ s_{l_e l_2 m} &= i^l (2l + 1) \sqrt{\frac{2l_e + 1}{2l_2 + 1}} \langle l, l_e, 0, m | l_2, m \rangle \langle l, l_e, 0, 0 | l_2, 0 \rangle. \end{aligned} \quad (30)$$

The line-width of an excited state l_e, n_e is approximately given by

$$R_{l_e n_e} = \sum_{ln} (1 - N_{ln}) SU_{ln l_e n_e} + N_{ln} SV_{ln l_e n_e}. \quad (31)$$

The matrix SV is obtained from the matrix SU by a substitution $u \rightarrow v$. As the eigenfunction $u_{ln}(r)$ evolves in the process of laser cooling it is more efficient to express the matrix (jWu) through a static matrix (jWW) build out of harmonic oscillator modes

$$(jWW)_{l_e n_e l_1 n_1} = \int r^2 dr j_l(kr) W_{l_e n_e}(r) W_{l_1 n_1}(r). \quad (32)$$

The static matrices s, S, jWW were prepared once, and stored then on a disk. They facilitate calculation of the temperature dependent matrices SU, su, R every time the wave functions u, v are updated.